

Intersection theorems for finite sets

Dhruv Mubayi

Department of Mathematics, Statistics and Computer Science
University of Illinois
Chicago

Robin Thomas 50

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

- $ax \leq 50$

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

- $ax \leq 50$
- $ay \leq 100$

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

- $ax \leq 50$
- $ay \leq 100$
- $bx \leq 100$

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

- $ax \leq 50$
- $ay \leq 100$
- $bx \leq 100$
- $by \leq 100$

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

- $ax \leq 50$
- $ay \leq 100$
- $bx \leq 100$
- $by \leq 100$

Prove that

$$(a + b)(x + y) \leq$$

A Puzzle

Suppose that a, b, x, y are positive real numbers such that

- $ax \leq 50$
- $ay \leq 100$
- $bx \leq 100$
- $by \leq 100$

Prove that

$$(a + b)(x + y) \leq 300.$$

The Frankl-Rödl theorem

Let M be a set. A family of sets \mathcal{A} is M -intersecting if
 $|A \cap B| \in M$ for every $A, B \in \mathcal{A}$

The Frankl-Rödl theorem

Let M be a set. A family of sets \mathcal{A} is M -intersecting if

$$|A \cap B| \in M \text{ for every } A, B \in \mathcal{A}$$

General Problem of Extremal Set Theory:

Given $\mathcal{A} \subset 2^{[n]}$ and $M \subset [n] := \{0, 1, \dots, n\}$, what is $\max |\mathcal{A}|$?

The Frankl-Rödl theorem

Let M be a set. A family of sets \mathcal{A} is M -intersecting if

$$|A \cap B| \in M \text{ for every } A, B \in \mathcal{A}$$

General Problem of Extremal Set Theory:

Given $\mathcal{A} \subset 2^{[n]}$ and $M \subset [n] := \{0, 1, \dots, n\}$, what is $\max |\mathcal{A}|$?

As M gets larger, $\max |\mathcal{A}|$ gets larger.

The Frankl-Rödl theorem

Let M be a set. A family of sets \mathcal{A} is M -intersecting if

$$|A \cap B| \in M \text{ for every } A, B \in \mathcal{A}$$

General Problem of Extremal Set Theory:

Given $\mathcal{A} \subset 2^{[n]}$ and $M \subset [n] := \{0, 1, \dots, n\}$, what is $\max |\mathcal{A}|$?

As M gets larger, $\max |\mathcal{A}|$ gets larger.

What if M misses only one number?

The Frankl-Rödl theorem

Let M be a set. A family of sets \mathcal{A} is M -intersecting if

$$|A \cap B| \in M \text{ for every } A, B \in \mathcal{A}$$

General Problem of Extremal Set Theory:

Given $\mathcal{A} \subset 2^{[n]}$ and $M \subset [n] := \{0, 1, \dots, n\}$, what is $\max |\mathcal{A}|$?

As M gets larger, $\max |\mathcal{A}|$ gets larger.

What if M misses only one number?

Theorem (Frankl-Rödl (1987), \$250 problem of Erdős)

Suppose that $\mathcal{A} \subset 2^{[n]}$ and $|A \cap B| \neq n/4$ for all $A, B \in \mathcal{A}$, and $n > n_0$. Then

$$|\mathcal{A}| < (1.99)^n.$$

Coding theory

- Q is an alphabet
- $q = |Q|$
- $\mathcal{C} \subset Q^n$ is a code

- Q is an alphabet
- $q = |Q|$
- $\mathcal{C} \subset Q^n$ is a code
- Hamming distance between codewords $C = (c_1, \dots, c_n)$ and $D = (d_1, \dots, d_n)$ is

$$d(C, D) := |\{i : c_i \neq d_i\}|$$

- Q is an alphabet
- $q = |Q|$
- $\mathcal{C} \subset Q^n$ is a code
- Hamming distance between codewords $C = (c_1, \dots, c_n)$ and $D = (d_1, \dots, d_n)$ is

$$d(C, D) := |\{i : c_i \neq d_i\}|$$

- $d(\mathcal{C}) = \{d(C, D) : C, D \in \mathcal{C}, C \neq D\}$

Problem. Find upper and lower bounds for $\max |\mathcal{C}|$ given $d(\mathcal{C})$.

Problem. Find upper and lower bounds for $\max |\mathcal{C}|$ given $d(\mathcal{C})$.

Theorem (Blokhuis, Frankl (1984))

Suppose that p is prime and $d(\mathcal{C})$ is covered by t nonzero residue classes mod p . Then

$$|\mathcal{C}| \leq \sum_{i=0}^t (q-1)^{n-i} \binom{n}{i} = q^n \sum_{i=0}^t (1-1/q)^{n-i} (1/q)^i \binom{n}{i}.$$

Problem. Find upper and lower bounds for $\max |\mathcal{C}|$ given $d(\mathcal{C})$.

Theorem (Blokhuis, Frankl (1984))

Suppose that p is prime and $d(\mathcal{C})$ is covered by t nonzero residue classes mod p . Then

$$|\mathcal{C}| \leq \sum_{i=0}^t (q-1)^{n-i} \binom{n}{i} = q^n \sum_{i=0}^t (1-1/q)^{n-i} (1/q)^i \binom{n}{i}.$$

If $t > n/q$, then concentration of the binomial distribution shows that the bound above is $q^{(1-o(1))n}$, which is rather weak.

Problem. Find upper and lower bounds for $\max |\mathcal{C}|$ given $d(\mathcal{C})$.

Theorem (Blokhuis, Frankl (1984))

Suppose that p is prime and $d(\mathcal{C})$ is covered by t nonzero residue classes mod p . Then

$$|\mathcal{C}| \leq \sum_{i=0}^t (q-1)^{n-i} \binom{n}{i} = q^n \sum_{i=0}^t (1-1/q)^{n-i} (1/q)^i \binom{n}{i}.$$

If $t > n/q$, then concentration of the binomial distribution shows that the bound above is $q^{(1-o(1))n}$, which is rather weak.

Theorem (Frankl-Rödl (1987))

Let $0 < \delta < 1/2$ and $\delta n < d < (1 - \delta)n$, and d is even if $q = 2$. If $d \notin d(\mathcal{C})$, then $|\mathcal{C}| < (q - \varepsilon)^n$, where $\varepsilon = \varepsilon(\delta, q) > 0$.

Borsuk's Problem - decreasing the diameter

Borsuk's Problem - decreasing the diameter

The Diameter of a set $S \subset \mathbb{R}^n$ is $\sup_{x,y \in S} \text{dist}(x,y)$

Borsuk's Problem - decreasing the diameter

The Diameter of a set $S \subset R^n$ is $\sup_{x,y \in S} \text{dist}(x,y)$

Conjecture

Every bounded $S \subset R^d$ can be partitioned into $d + 1$ sets S_1, \dots, S_{d+1} of smaller diameter.

Borsuk's Problem - decreasing the diameter

The Diameter of a set $S \subset \mathbb{R}^n$ is $\sup_{x,y \in S} \text{dist}(x,y)$

Conjecture

Every bounded $S \subset \mathbb{R}^d$ can be partitioned into $d + 1$ sets S_1, \dots, S_{d+1} of smaller diameter.

If true, then sharp by letting S be the vertices of a regular simplex, for example,

$$S = \{e_1, \dots, e_d, v\}$$

where e_i is the unit vector with 1 in position i , and

$$v = \frac{1 - \sqrt{n+1}}{n}(1, \dots, 1).$$

- Borsuk (1932) $d = 2$
- Eggleston (1955) $d = 3$
- Hadwiger (1946) for all d if S is smooth and convex
- Riesling (1971) for all d if S is centrally symmetric
- Dekster (1995) for all d if S is a body of revolution
- Schramm (1988) number of pieces is at most $(\sqrt{3/2} + \epsilon)^d$, for all $\epsilon > 0$ and $d > d(\epsilon)$.

Counterexamples

Theorem (Kahn-Kalai (1993))

For large d , there exists a bounded $S \subset R^d$ such that every partition of S into pieces of smaller diameter has at least $(1.2)^{\sqrt{d}}$ parts. In particular, Borsuk's conjecture fails for $d = 1325$ and each $d > 2014$.

Counterexamples

Theorem (Kahn-Kalai (1993))

For large d , there exists a bounded $S \subset R^d$ such that every partition of S into pieces of smaller diameter has at least $(1.2)^{\sqrt{d}}$ parts. In particular, Borsuk's conjecture fails for $d = 1325$ and each $d > 2014$.

Conjecture

There exists $c > 1$ such that for all d , there exists a bounded $S \subset R^d$ such that every partition of S into pieces of smaller diameter has at least c^d parts.

More Geometry

How many vectors of the cube in R^d can be pairwise non-orthogonal?

More Geometry

How many vectors of the cube in R^d can be pairwise non-orthogonal?

Conjecture (Larman-Rogers (1972))

Suppose that $d = 4n$. Does every set of $2^d/d^2 \pm 1$ vectors in R^d contain a pair of orthogonal vectors?

More Geometry

How many vectors of the cube in R^d can be pairwise non-orthogonal?

Conjecture (Larman-Rogers (1972))

Suppose that $d = 4n$. Does every set of $2^d/d^2 \pm 1$ vectors in R^d contain a pair of orthogonal vectors?

Theorem (Frankl-Rödl (1987))

Given $r \geq 2$ and $n = d/4 \geq r$, there exists $\varepsilon = \varepsilon(r) > 0$ such that every set of more than $(2 - \varepsilon)^d \pm 1$ vectors in R^d contains r pairwise orthogonal vectors.

A weak delta system is a collection of sets A_1, \dots, A_r such that

$$|A_i \cap A_j| = |A_1 \cap A_2|$$

for $1 \leq i < j \leq r$.

A weak delta system is a collection of sets A_1, \dots, A_r such that

$$|A_i \cap A_j| = |A_1 \cap A_2|$$

for $1 \leq i < j \leq r$.

Conjecture (Erdős-Szemerédi (1978))

For every $\varepsilon > 0$, there is $n_0 = n_0(\varepsilon)$ such that if $n > n_0$ and $\mathcal{A} \subset 2^{[n]}$ with $|\mathcal{A}| > (2 - \varepsilon)^n$, then \mathcal{A} contains a weak delta system of size 3.

Theorem (Frankl-Rödl (1987))

Fix $r \geq 3$. Then there are $\eta = \eta(r)$ and $\varepsilon = \varepsilon(r)$ such that if $t = (1/4 \pm \eta)n$ and $\mathcal{A} \subset 2^{[n]}$ with $|\mathcal{A}| > (2 - \varepsilon)^n$, then there are $A_1, \dots, A_r \in \mathcal{A}$ with

$$|A_i \cap A_j| = t$$

for $1 \leq i < j \leq r$.

Theorem (Frankl-Rödl (1987))

Fix $r \geq 3$. Then there are $\eta = \eta(r)$ and $\varepsilon = \varepsilon(r)$ such that if $t = (1/4 \pm \eta)n$ and $\mathcal{A} \subset 2^{[n]}$ with $|\mathcal{A}| > (2 - \varepsilon)^n$, then there are $A_1, \dots, A_r \in \mathcal{A}$ with

$$|A_i \cap A_j| = t$$

for $1 \leq i < j \leq r$.

Conjecture (Erdős-Szemerédi (1978))

There exists $\varepsilon > 0$ such that if n is sufficiently large and $\mathcal{A} \subset 2^{[n]}$ with $|\mathcal{A}| > (2 - \varepsilon)^n$, then \mathcal{A} contains a delta system (not weak!) of size 3.

Theorem (Frankl-Rödl (1987))

Fix $r \geq 3$. Then there are $\eta = \eta(r)$ and $\varepsilon = \varepsilon(r)$ such that if $t = (1/4 \pm \eta)n$ and $\mathcal{A} \subset 2^{[n]}$ with $|\mathcal{A}| > (2 - \varepsilon)^n$, then there are $A_1, \dots, A_r \in \mathcal{A}$ with

$$|A_i \cap A_j| = t$$

for $1 \leq i < j \leq r$.

Conjecture (Erdős-Szemerédi (1978))

There exists $\varepsilon > 0$ such that if n is sufficiently large and $\mathcal{A} \subset 2^{[n]}$ with $|\mathcal{A}| > (2 - \varepsilon)^n$, then \mathcal{A} contains a delta system (not weak!) of size 3.

Recent work of Alon-Shpilka-Umans gives connections between this conjecture and algorithms for Matrix multiplication

- Communication Complexity (Sgall 1999)
- Quantum Computing (Buhrman-Cleve-Wigderson 1998)
- Semidefinite Programming (Goemans-Kleinberg 1998, Hatami-Magen-Markakis 2009)

The Frankl-Rödl theorem

The Frankl-Rödl theorem

Theorem (Frankl-Rödl (1987))

Let $0 < \eta < 1/2$ and $\eta n < t < (1/2 - \eta)n$. There is $\varepsilon_0 = \varepsilon_0(\eta)$ such that if $\mathcal{A} \subset 2^{[n]}$ and $|A \cap B| \neq t$ for all $A, B \in \mathcal{A}$, then

$$|\mathcal{A}| < (2 - \varepsilon_0)^n.$$

The Frankl-Rödl theorem

Theorem (Frankl-Rödl (1987))

Let $0 < \eta < 1/2$ and $\eta n < t < (1/2 - \eta)n$. There is $\varepsilon_0 = \varepsilon_0(\eta)$ such that if $\mathcal{A} \subset 2^{[n]}$ and $|A \cap B| \neq t$ for all $A, B \in \mathcal{A}$, then

$$|\mathcal{A}| < (2 - \varepsilon_0)^n.$$

How big is ε_0 ?

The Frankl-Rödl theorem

Theorem (Frankl-Rödl (1987))

Let $0 < \eta < 1/2$ and $\eta n < t < (1/2 - \eta)n$. There is $\varepsilon_0 = \varepsilon_0(\eta)$ such that if $\mathcal{A} \subset 2^{[n]}$ and $|A \cap B| \neq t$ for all $A, B \in \mathcal{A}$, then

$$|\mathcal{A}| < (2 - \varepsilon_0)^n.$$

How big is ε_0 ?

Frankl-Rödl show it is about $(t/n)^2/2$.

The optimal ε_0

The binary entropy function is

$$H(x) = -x \log_2 x - (1 - x) \log_2(1 - x).$$

The optimal ε_0

The binary entropy function is

$$H(x) = -x \log_2 x - (1 - x) \log_2 (1 - x).$$

Conjecture (M-Rödl)

Let $0 < \eta < 1/3$, $\eta n < t < n/3$, and $\mathcal{A} \subset 2^{[n]}$ with $|A \cap B| \neq t$ for all $A, B \in \mathcal{A}$. Then

$$|\mathcal{A}| \leq \binom{n}{(n+t)/2} 2^{o(n)} = 2^{H(\frac{1}{2} + \frac{t}{2n})n + o(n)}.$$

The optimal ε_0

The binary entropy function is

$$H(x) = -x \log_2 x - (1 - x) \log_2(1 - x).$$

Conjecture (M-Rödl)

Let $0 < \eta < 1/3$, $\eta n < t < n/3$, and $\mathcal{A} \subset 2^{[n]}$ with $|A \cap B| \neq t$ for all $A, B \in \mathcal{A}$. Then

$$|\mathcal{A}| \leq \binom{n}{(n+t)/2} 2^{o(n)} = 2^{H(\frac{1}{2} + \frac{t}{2n})n + o(n)}.$$

If true, the conjecture is (asymptotically) sharp via $\mathcal{A} = \binom{[n]}{>_{(n+t)/2}}$.

The optimal ε_0

The binary entropy function is

$$H(x) = -x \log_2 x - (1-x) \log_2 (1-x).$$

Conjecture (M-Rödl)

Let $0 < \eta < 1/3$, $\eta n < t < n/3$, and $\mathcal{A} \subset 2^{[n]}$ with $|A \cap B| \neq t$ for all $A, B \in \mathcal{A}$. Then

$$|\mathcal{A}| \leq \binom{n}{(n+t)/2} 2^{o(n)} = 2^{H(\frac{1}{2} + \frac{t}{2n})n + o(n)}.$$

If true, the conjecture is (asymptotically) sharp via $\mathcal{A} = \binom{[n]}{>_{(n+t)/2}}$.

For $n/3 < t < (1/2 - \eta)n$, the construction $\mathcal{A} = \binom{[n]}{t}$ is better, and we conjecture it is optimal.

Forbidding a small interval

Forbidding a small interval

Theorem (M-Rödl)

Let $0 < \varepsilon < 1/5$ be fixed, $n > n_0(\varepsilon)$, $\varepsilon n < t < n/5$ and $\mathcal{A} \subset 2^{[n]}$. Suppose that

$$|A \cap B| \notin (t, t + n^{0.525})$$

for all $A, B \in \mathcal{A}$. Then

$$|\mathcal{A}| < n \binom{n}{(n+t)/2}.$$

Forbidding a small interval

Theorem (M-Rödl)

Let $0 < \varepsilon < 1/5$ be fixed, $n > n_0(\varepsilon)$, $\varepsilon n < t < n/5$ and $\mathcal{A} \subset 2^{[n]}$. Suppose that

$$|A \cap B| \notin (t, t + n^{0.525})$$

for all $A, B \in \mathcal{A}$. Then

$$|\mathcal{A}| < n \binom{n}{(n+t)/2}.$$

- The constant 0.525 is a consequence of the result of Baker-Harman-Pintz that there is a prime in every interval $(s - s^{0.525}, s)$ as long as s is sufficiently large.

Forbidding a small interval

Theorem (M-Rödl)

Let $0 < \varepsilon < 1/5$ be fixed, $n > n_0(\varepsilon)$, $\varepsilon n < t < n/5$ and $\mathcal{A} \subset 2^{[n]}$. Suppose that

$$|A \cap B| \notin (t, t + n^{0.525})$$

for all $A, B \in \mathcal{A}$. Then

$$|\mathcal{A}| < n \binom{n}{(n+t)/2}.$$

- The constant 0.525 is a consequence of the result of Baker-Harman-Pintz that there is a prime in every interval $(s - s^{0.525}, s)$ as long as s is sufficiently large.
- If we assume the Riemann Hypothesis, then 0.525 could be improved to $1/2 + o(1)$ using a result of Cramér.

More restricted intersections

Question. Can the upper bound for M -intersecting families be improved for more restrictive M ?

More restricted intersections

Question. Can the upper bound for M -intersecting families be improved for more restrictive M ?

Theorem (Berlekamp (1965), Graver (1975))

Suppose that $\mathcal{A} \subset 2^{[n]}$ is M -intersecting, where $M = \{0, 2, 4, \dots\}$. In other words, $|A \cap B|$ is even for all $A, B \in \mathcal{A}$. Then $|\mathcal{A}| \leq 2^{\lfloor n/2 \rfloor} + 1$.

More restricted intersections

Question. Can the upper bound for M -intersecting families be improved for more restrictive M ?

Theorem (Berlekamp (1965), Graver (1975))

Suppose that $\mathcal{A} \subset 2^{[n]}$ is M -intersecting, where $M = \{0, 2, 4, \dots\}$. In other words, $|A \cap B|$ is even for all $A, B \in \mathcal{A}$. Then $|\mathcal{A}| \leq 2^{\lfloor n/2 \rfloor} + 1$.

Eventown Theorem

Suppose that $\mathcal{A} \subset 2^{[n]}$ such that

- $|A|$ is even for every $A \in \mathcal{A}$
- $|A \cap B|$ is even for every $A, B \in \mathcal{A}$

Then $|\mathcal{A}| \leq 2^{\lfloor n/2 \rfloor}$.

A Proof

Proof of Eventown:

- To each $A \in \mathcal{A}$, associate its incidence vector $v_A = (v_1, \dots, v_n)$ where

$$v_i = \begin{cases} 1 & i \in A \\ 0 & i \notin A \end{cases}$$

A Proof

Proof of Eventown:

- To each $A \in \mathcal{A}$, associate its incidence vector $v_A = (v_1, \dots, v_n)$ where

$$v_i = \begin{cases} 1 & i \in A \\ 0 & i \notin A \end{cases}$$

- Let S be the subspace of F_2^n spanned by $\{v_A\}_{A \in \mathcal{A}}$.

A Proof

Proof of Eventown:

- To each $A \in \mathcal{A}$, associate its incidence vector $v_A = (v_1, \dots, v_n)$ where

$$v_i = \begin{cases} 1 & i \in A \\ 0 & i \notin A \end{cases}$$

- Let S be the subspace of F_2^n spanned by $\{v_A\}_{A \in \mathcal{A}}$.
 - S is totally isotropic (meaning $x \cdot y = 0$ for $x, y \in S$)
 - $\dim(S) \leq \lfloor n/2 \rfloor$

A Proof

Proof of Eventown:

- To each $A \in \mathcal{A}$, associate its incidence vector $v_A = (v_1, \dots, v_n)$ where

$$v_i = \begin{cases} 1 & i \in A \\ 0 & i \notin A \end{cases}$$

- Let S be the subspace of F_2^n spanned by $\{v_A\}_{A \in \mathcal{A}}$.
 - S is totally isotropic (meaning $x \cdot y = 0$ for $x, y \in S$)
 - $\dim(S) \leq \lfloor n/2 \rfloor$
- So $|\mathcal{A}| \leq |S| \leq 2^{\lfloor n/2 \rfloor} = (1.4142\dots)^n$

Between the extremes

Frankl-Rödl: $M = [n] \setminus \{n/4\}$ — $|\mathcal{A}| < (1.99)^n$

Between the extremes

Frankl-Rödl: $M = [n] \setminus \{n/4\}$ — $|\mathcal{A}| < (1.99)^n$

Eventown: $M = \{0, 2, \dots\}$ — $|\mathcal{A}| < (1.4142\dots)^n$

Between the extremes

Frankl-Rödl: $M = [n] \setminus \{n/4\}$ — $|\mathcal{A}| < (1.99)^n$

Eventown: $M = \{0, 2, \dots\}$ — $|\mathcal{A}| < (1.4142\dots)^n$

What about M that are in between these two extremes?

Between the extremes

Frankl-Rödl: $M = [n] \setminus \{n/4\}$ — $|\mathcal{A}| < (1.99)^n$

Eventown: $M = \{0, 2, \dots\}$ — $|\mathcal{A}| < (1.4142\dots)^n$

What about M that are in between these two extremes?

Definition

The length $\ell(M)$ of a set M is the maximum number of consecutive integers contained in M .

Between the extremes

Frankl-Rödl: $M = [n] \setminus \{n/4\}$ — $|\mathcal{A}| < (1.99)^n$

Eventown: $M = \{0, 2, \dots\}$ — $|\mathcal{A}| < (1.4142\dots)^n$

What about M that are in between these two extremes?

Definition

The length $\ell(M)$ of a set M is the maximum number of consecutive integers contained in M .

$\ell(M) \leq \ell$ if and only if \overline{M} is $(\ell + 1)$ -syndetic.

Bounds for small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 1.622^n \times 10^{2\ell+5}.$$

Bounds for small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 1.622^n \times 10^{2\ell+5}.$$

- The result is nontrivial as long as, $\ell < n/10$ or so

Bounds for small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 1.622^n \times 10^{2\ell+5}.$$

- The result is nontrivial as long as, $\ell < n/10$ or so
- For example, if $[n] \setminus M = \{0, n/10^4, 2n/10^4, \dots\}$, then

$$|\mathcal{A}| < 1.63^n.$$

Bounds for small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 1.622^n \times 10^{2\ell+5}.$$

- The result is nontrivial as long as, $\ell < n/10$ or so
- For example, if $[n] \setminus M = \{0, n/10^4, 2n/10^4, \dots\}$, then

$$|\mathcal{A}| < 1.63^n.$$

- The 1.622 is probably not sharp, just a result of the proof

Bounds for very small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 2^{n/2 + \ell \log^2 n}.$$

Bounds for very small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 2^{n/2 + \ell \log^2 n}.$$

- For $\ell = o(n / \log^2 n)$, this bound better than the first one; it is

$$|\mathcal{A}| < 2^{n/2 + o(n)}.$$

Bounds for very small $\ell(M)$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $\mathcal{A} \subset 2^{[n]}$ is an M -intersecting family. Then

$$|\mathcal{A}| < 2^{n/2 + \ell \log^2 n}.$$

- For $\ell = o(n / \log^2 n)$, this bound better than the first one; it is

$$|\mathcal{A}| < 2^{n/2 + o(n)}.$$

- This is the first non-linear-algebraic proof of an asymptotic version of the Eventown Theorem; it applies in more general scenarios though doesn't give bounds as precise as $2^{n/2}$.

- Prove the result for pairs of families $(\mathcal{A}, \mathcal{B})$. This facilitates an induction argument

- Prove the result for pairs of families $(\mathcal{A}, \mathcal{B})$. This facilitates an induction argument
- $(\mathcal{A}, \mathcal{B})$ is M -intersecting if

$$|A \cap B| \in M$$

for all $A \in \mathcal{A}$ and $B \in \mathcal{B}$

- Prove the result for pairs of families $(\mathcal{A}, \mathcal{B})$. This facilitates an induction argument
- $(\mathcal{A}, \mathcal{B})$ is M -intersecting if

$$|A \cap B| \in M$$

for all $A \in \mathcal{A}$ and $B \in \mathcal{B}$

Theorem (M-Rödl)

Let $M \subset [n]$ with $\ell(M) = \ell$. Suppose that $(\mathcal{A}, \mathcal{B})$ is an M -intersecting pair of families in $2^{[n]}$. Then

$$|\mathcal{A}||\mathcal{B}| < \min \left\{ 2.631^n \times 10^{4\ell+10}, \quad 2^{n+2\ell \log^2 n} \right\}.$$

Height functions

Height functions

Definition (Sgall)

Say that a function $h : N^{<\infty} \rightarrow N$ is a *height function* if the following four properties hold:

Height functions

Definition (Sgall)

Say that a function $h : N^{<\infty} \rightarrow N$ is a *height function* if the following four properties hold:

(A1) $h(L) = 0$ if and only if $L = \emptyset$,

Height functions

Definition (Sgall)

Say that a function $h : N^{<\infty} \rightarrow N$ is a *height function* if the following four properties hold:

(A1) $h(L) = 0$ if and only if $L = \emptyset$,

(A2) if $L' \subset L$, then $h(L') \leq h(L)$,

Height functions

Definition (Sgall)

Say that a function $h : N^{<\infty} \rightarrow N$ is a *height function* if the following four properties hold:

- (A1) $h(L) = 0$ if and only if $L = \emptyset$,
- (A2) if $L' \subset L$, then $h(L') \leq h(L)$,
- (A3) if $L' \subset L - 1$, then $h(L') \leq h(L)$,

Height functions

Definition (Sgall)

Say that a function $h : N^{<\infty} \rightarrow N$ is a *height function* if the following four properties hold:

(A1) $h(L) = 0$ if and only if $L = \emptyset$,

(A2) if $L' \subset L$, then $h(L') \leq h(L)$,

(A3) if $L' \subset L - 1$, then $h(L') \leq h(L)$,

(A4) if $h(L), h(L') \leq s$, then either

$$h(L' \cap L) \leq s - 1 \quad \text{or} \quad h(L' \cap (L - 1)) \leq s - 1.$$

Sgall's theorem

Theorem (Sgall (1999))

Suppose that $(\mathcal{A}, \mathcal{B})$ is an M -intersecting pair of families in $2^{[n]}$ and $h(M) \leq s \leq n+1$. Then

$$|\mathcal{A}||\mathcal{B}| \leq 2^{n+s-1} \binom{n}{s-1}.$$

Sgall's theorem

Theorem (Sgall (1999))

Suppose that $(\mathcal{A}, \mathcal{B})$ is an M -intersecting pair of families in $2^{[n]}$ and $h(M) \leq s \leq n + 1$. Then

$$|\mathcal{A}||\mathcal{B}| \leq 2^{n+s-1} \binom{n}{s-1}.$$

Theorem (M-Rödl)

There exists a height function h such that for every $M \subset [n]$,

$$h(M) \leq 1 + 2\ell(M) \log n.$$

Applying this bound in Sgall's Theorem yields $|\mathcal{A}||\mathcal{B}| < 2^{n+2\ell \log^2 n}$.

The Height function

- $h(\emptyset) = 0$

The Height function

- $h(\emptyset) = 0$
- Suppose that $L \neq \emptyset$ and h has been defined on all sets of size less than $|L|$

The Height function

- $h(\emptyset) = 0$
- Suppose that $L \neq \emptyset$ and h has been defined on all sets of size less than $|L|$
- $T(L) = \{M : M \notin \{L, L+1\} \text{ and } 0 < |M| \leq |L|\}$

The Height function

- $h(\emptyset) = 0$
- Suppose that $L \neq \emptyset$ and h has been defined on all sets of size less than $|L|$
- $T(L) = \{M : M \not\subseteq \{L, L+1\} \text{ and } 0 < |M| \leq |L|\}$
- $A = h(L \cap (L+1))$

The Height function

- $h(\emptyset) = 0$
- Suppose that $L \neq \emptyset$ and h has been defined on all sets of size less than $|L|$
- $T(L) = \{M : M \not\subseteq \{L, L+1\} \text{ and } 0 < |M| \leq |L|\}$
- $A = h(L \cap (L+1))$
- $B = \max_{M \in T(L)} \min\{h(L \cap M), h(L \cap (M-1))\}$

The Height function

- $h(\emptyset) = 0$
- Suppose that $L \neq \emptyset$ and h has been defined on all sets of size less than $|L|$
- $T(L) = \{M : M \not\subseteq \{L, L+1\} \text{ and } 0 < |M| \leq |L|\}$
- $A = h(L \cap (L+1))$
- $B = \max_{M \in T(L)} \min\{h(L \cap M), h(L \cap (M-1))\}$
- $h(L) = 1 + \max\{A, B\}$

Sgall's Lemma and the Puzzle

Lemma (Sgall)

Suppose that a, b, x, y, p, Q are positive real numbers such that

Sgall's Lemma and the Puzzle

Lemma (Sgall)

Suppose that a, b, x, y, p, Q are positive real numbers such that

- $ax \leq p$
- $ay \leq Q$
- $bx \leq Q$
- $by \leq Q$

Sgall's Lemma and the Puzzle

Lemma (Sgall)

Suppose that a, b, x, y, p, Q are positive real numbers such that

- $ax \leq p$
- $ay \leq Q$
- $bx \leq Q$
- $by \leq Q$

Then

$$(a + b)(x + y) \leq$$

Sgall's Lemma and the Puzzle

Lemma (Sgall)

Suppose that a, b, x, y, p, Q are positive real numbers such that

- $ax \leq p$
- $ay \leq Q$
- $bx \leq Q$
- $by \leq Q$

Then

$$(a + b)(x + y) \leq 2(p + Q).$$

Happy Birthday Robin!